

## Perturbation Spreading in Many-Particle Systems: A Random Walk Approach

V. Zaburdaev,<sup>1</sup> S. Denisov,<sup>2</sup> and P. Hänggi<sup>2</sup>

<sup>1</sup>*School of Engineering and Applied Science, Harvard University, 29 Oxford Street, Cambridge, Massachusetts 02138, USA*

<sup>2</sup>*Institute of Physics, University of Augsburg, Universitätsstrasse 1, D-86159 Augsburg, Germany*

(Received 7 January 2011; published 3 May 2011)

The propagation of an initially localized perturbation via an interacting many-particle Hamiltonian dynamics is investigated. We argue that the propagation of the perturbation can be captured by the use of a continuous-time random walk where a single particle is traveling through an active, fluctuating medium. Employing two archetype ergodic many-particle systems, namely, (i) a hard-point gas composed of two unequal masses and (ii) a Fermi-Pasta-Ulam chain, we demonstrate that the corresponding perturbation profiles coincide with the diffusion profiles of the single-particle Lévy walk approach. The parameters of the random walk can be related through elementary algebraic expressions to the physical parameters of the corresponding test many-body systems.

DOI: [10.1103/PhysRevLett.106.180601](https://doi.org/10.1103/PhysRevLett.106.180601)

PACS numbers: 05.40.Fb, 05.45.Jn, 45.50.Jf

The transport properties of many-particle systems are of salient interest in diverse contexts, ranging from foundations of thermodynamics to the transduction of information on the nanoscale. The collective evolution of  $N \gg 1$  interacting particles creates a dynamical “tissue,” whose properties depend not only on the Hamiltonian of the system, but also on the state of the system itself. An objective of primary interest is how the system responds to the perturbation that locally affects its dynamics. The answer then provides direct insight into collective energy, correlation, and information transport in extended nondissipative media [1–7].

Consider the situation of a many-particle system at microcanonical equilibrium, when at the initial time  $t = 0$  one of the particles receives some external local perturbation. The system gains a small amount of perturbation energy, which is conserved due to Hamiltonian evolution of the system. However, the perturbation does spread as the perturbation energy is shared by a constantly growing number of particles. One of the main features of the spreading is the finite velocity of perturbation propagation,  $v_0 < \infty$  [4,6,7]. Therefore, similar to relativistic diffusion theories [8], an effective “light cone” occurs [9], such that at the given time  $t$  the perturbation is almost completely confined to the interval  $[-v_0t, v_0t]$  [10]. The fundamental fact of the cone’s existence assumes the characteristics of a mathematical existence theorem [9,11]; its strength therefore is the generality when dealing with many-particle systems in rather arbitrary situations. A pronounced weakness of this mathematical approach, however, becomes evident whenever one attempts to implement the theory for a particular system to obtain qualitative results on an analytical level of description. Therefore, more applied approaches, like the diffusion formalism, might become of beneficial use. Indeed it is appealing to consider the said perturbation spreading as a certain (not yet known) one-dimensional diffusion process, and to

quantify it with the mean square displacement and other related attributes [1,5–7]. As it is known, however, it is impossible to describe this process by any known macroscopic, norm-preserving normal diffusion equation. Conventional diffusion equations knowingly lead to infinite propagation speeds [8] and therefore are incompatible with the existence of a causal cone. Thus the perturbation kinetics should be ultimately considered on the microscopic level corresponding to the random walk approach.

With this Letter we employ the microscopic single-particle random walk process in order to evaluate the evolution of perturbations in one-dimensional, ergodic many-particle systems. In doing so we address the two challenges: (i) How is the perturbation distributed within the cone, and (ii) what are the shapes of the cone fronts? By using two renowned many-particle chains we demonstrate that the random walk model accurately describes the perturbation spreading in these interacting many-body systems. We show that the walker-media interactions are responsible for the observed shape of the causal cone and predict the universal scalings for the perturbation profile and its corresponding fronts.

*Model setup.*—The continuous-time random walk (CTRW) formalism [12] has found applications to a wide range of phenomena, ranging from financial markets dynamics [13] to single molecule spectroscopy [14]. Here, we use one specific CTRW model [15,16], where a walker moves ballistically in between successive “turning points.” During a single flight event, the walker travels at constant speed  $v_0$ , and at the turning points it randomly changes the direction of motion (see the particle flying above the slab in Fig. 1). Hence, the velocity probability density function (PDF) reads  $h(v) = [\delta(v - v_0) + \delta(v + v_0)]/2$ . The flight times,  $\tau_n$ , are independent and identically distributed random variables drawn from a PDF  $\psi(\tau)$  that is described by a power law [5,16–19],




FIG. 1 (color online). The standard continuous-time random walk (above the slab) and random walk through an active medium (in the slab). The first walker performs flights with constant velocity  $\pm v_0$ , while the second one is subjected to constant scattering, so that its velocity fluctuates. The duration of single flight,  $\tau_n$ , is a random variable drawn from the probability density function  $\psi(\tau)$ .

$$\psi(\tau) \propto (\tau/\tau_0)^{-\gamma-1}, \quad (1)$$

where  $\tau_0$  is a characteristic time scale, and  $1 < \gamma \leq 2$ . This choice guarantees a finite average flight time,  $\langle \tau \rangle = \int_0^\infty \tau \psi(\tau) d\tau$ , and provides access to different diffusion regimes with the scaling of the mean squared displacement  $\sigma^2(t) \propto t^{3-\gamma}$ . The corresponding Lévy walk (LW) approach [16] has been successfully used for the description of diffusion of particles in chaotic systems [17–19], tracers in turbulent flows [20], or ultracold atoms in optical potentials [21]. For the standard LW process, the PDF of finding a particle in  $x$  at the time  $t$ ,  $P(x, t)$ , provided it was initially localized at  $x = 0$ , exhibits a sharp cutoff marked by the ballistic peaks at  $|x| = v_0 t$  [19].

A first step towards the implementation of the LW formalism for the description of the perturbation dynamics in many-particle systems has been attempted in [5]. Although this approach provided an adequate description for the perturbation spreading process, it failed to capture the dynamics of the cone fronts. The observed ballistic humps displayed smooth, Gaussian-like profiles, with a scaling that was incompatible with the scaling behavior of deltalike peaks of the LW propagator [18].

In order to resolve this issue, we here extend the conventional CTRW setup by assuming that the walker performs a random walk through an active medium. Conceptually, it means that while moving, the walker interacts with the surrounding medium (the slab in Fig. 1). This interaction causes fluctuations of the walker's velocity (note the noisy trajectory of the particle moving inside the slab in Fig. 1). The term “active” means that the medium is not solely dissipative and the particle not only continuously loses but also gains energy from its environment. Both processes are in balance, yielding unbiased fluctuations of the walker's velocity around  $v_0$ .

*Model dynamics.*—We start out by considering a single flight event. The position of the walker is defined by a Langevin equation  $\dot{x} = v_0 + \xi(t)$ , where  $\xi(t)$  is a delta-correlated Gaussian process of vanishing mean and finite intensity  $D_v$ , i.e.,  $\langle \xi(t)\xi(s) \rangle = D_v \delta(t-s)$ . This constitutes a well-known biased Wiener process with drift  $v_0$  [22]. After an integration over a time interval  $\tau$  we obtain




FIG. 2 (color online). Rescaled propagators of Lévy walk process with exponent  $\gamma = 5/3$  for different times,  $t = 100, 300, 600$ . The dashed line depicts the propagator for the standard Lévy walk with the constant flight velocity  $v_0 = 1$ . For the simulations of the random walk in the active medium we employed the map (2) with  $D_v = 0.03$ . The inset shows the ballistic front regions after the scaling transformation (5), where  $\bar{x} = x - v_0 t$ .

$$x(t + \tau) = x(t) + v_0 \tau + w(\tau), \quad (2)$$

where the new stochastic variable  $w(\tau) = \int_t^{t+\tau} \xi(s) ds$  is characterized by the Gaussian PDF  $p(w, \tau)$  with the dispersion  $\sigma_w^2 = \langle [x(\tau) - v_0 \tau]^2 \rangle = D_v \tau$ . The propagator can be calculated numerically by sampling long enough a single-particle trajectory [23] (see Fig. 2).

To gain analytical insight into the generalized LW dynamics, we follow a standard reasoning [18,24] and derive the transport equations governing the evolution of the particle density,  $P(x, t)$  (see supplemental material [25]). Below we present the major results for the scaling properties of the central part of the density profile and provide the explicit expression describing the ballistic humps. The asymptotic analysis of the central part of the density profile reveals the scaling of the standard LW propagator [18], namely,

$$P(x, t') \simeq \frac{1}{K u^{1/\gamma}} P\left(\frac{x}{K u^{1/\gamma}}, t\right), \quad |x| \ll v t, \quad (3)$$

where  $K \propto \tau_0^{1-1/\gamma} v_0$  and  $u = t'/t$  (see Fig. 2).

The salient difference between the dynamics of our model and the standard Lévy walk becomes apparent in the regions of cone fronts. The ballistic humps are formed by the particles which were flying from  $t = 0$  to the observation time  $t$  (see supplemental material [25]):

$$P_{\text{hump}}(x, t) = \Phi(t)[p(x + v_0 t, t) + p(x - v_0 t, t)]/2. \quad (4)$$

Here,  $\Phi(t)$  denotes the probability of not changing the direction of flight during the time  $t$  and has a power-law behavior  $\Phi(t) \propto (t/\tau_0)^{1-\gamma}$  [18]. Consequently, the area under the ballistic humps (4) also scales as  $t^{1-\gamma}$ . During flights, the particles undergo random fluctuations caused by the flight's velocity variations. The flight length is proportional to  $t$ ; thus the dispersion of the Gaussian-like

humps grows as  $\sqrt{t}$ , and we arrive at the following scaling for the particles' density in the hump regions:

$$P_{\text{hump}}(\bar{x}, t') \simeq u^{-1/2} P_{\text{hump}}(\bar{x}/u^{\gamma-1/2}, t), \quad (5)$$

where  $u = t'/t$  and  $\bar{x} = x - v_0 t$  (see inset in Fig. 2). Note that this scaling distinctly differs from the scaling in Eq. (3).

*Ergodic many-body systems: Validation of the approach.*—Consider a many-particle system, with a Hamiltonian

$$H_{\text{total}}(\{x_i, p_i\}) = \sum_{i=1}^N H_i, \quad (6)$$

where  $H_i = H(x_i, x_{i-1}, x_{i+1}, p_i)$  is the energy attributed to the  $i$ th particle. At the time  $t = 0$  the system is locally affected by the perturbation. The initially localized perturbation energy,  $E_p$ , starts to spread, such that the distribution of the local excess energy  $\Delta E(i, t)$  of the  $i$  particle evolves in time [26], while keeping the perturbation energy constant; i.e.,  $\sum_{i=1}^N \Delta E(i, t) = E_p$ . The spreading can be quantified with a normalized probability distribution function  $\varrho(i, t) = \overline{\Delta E(i, t)}/E_p$ , where  $\overline{\dots}$  denotes a microcanonical average.

The main finding of this study is that the profiles of the spreading perturbation in many-particle Hamiltonian systems,  $\varrho(i, t)$ , in the corresponding asymptotic regime [27], are determined by the propagator  $P(x, t)$  of the generalized Lévy walk model, Eqs. (3) and (5). In order to validate this claim we use two archetype systems, namely, (i) a one-dimensional scattering dynamics of a hard-point gas composed of two unequal masses [28] and (ii) a Fermi-Pasta-Ulam (FPU)  $\beta$ -lattice dynamics [29].

We start with a hard-point gas, a many-body Hamiltonian system with an ergodic dynamics governed by the conservation of kinetic energy and momentum [28]. We use a chain of  $N = 1.6 \times 10^4$  pointlike particles with alternating masses,  $\dots m M m M \dots$ , of the length  $L = 1.6 \times 10^4$ , and periodic boundary conditions. Without loss of generality we set the mass ratio  $M/m = 2$ . The energy per particle,  $\varepsilon = \langle m_i v_i^2 \rangle / 2$ ,  $m_i = m$  or  $M$ , serves as a tunable parameter. Figure 3 depicts the evolution of the infinitesimal perturbation  $\varrho(i, t)$  [5]. The scaling ansatz (5) with the exponent  $\gamma = 5/3$  is beautifully validated (see the inset in Fig. 3(a)). We also found that the perturbation profiles for different values of energy per particle parameter  $\varepsilon$  are matched by assuming that the perturbation velocity and the fluctuation variance both scale as

$$v_0, D_v \propto \sqrt{\varepsilon}. \quad (7)$$

Consequently, the profile scales as

$$\varrho_\varepsilon(x, t) = \varrho_{\varepsilon'}(x, t/s'), \quad (8)$$

where  $s' = \sqrt{\varepsilon'/\varepsilon}$  (see Fig. 3(b)).




FIG. 3 (color online). (a) Rescaled perturbation profiles at times  $t = 1000, 2000, 4000$ , and  $6000$  (the width increases with time), for the hard-point gas dynamics with the energy per particle  $\varepsilon = 1$ . The scaling exponent is  $\gamma = 5/3$ . The inset, similar to the inset in Fig. 2, depicts the ballistic humps after the scaling transformation (5). (b) Profiles after the scaling transformation (8) for different energy per particle,  $\varepsilon$ , at time  $t = 1500/\sqrt{\varepsilon/\varepsilon_0}$ , with  $\varepsilon_0 = 1$ . Each profile is obtained by averaging over  $10^6$  realizations.

As our second test bed we use a FPU  $\beta$  chain dynamics [29], defined by the Hamiltonian (6) with  $H_i = \frac{1}{2} p_i^2 + \frac{1}{2} \times (x_{i+1} - x_i)^2 + \frac{\beta}{4} (x_{i+1} - x_i)^4$ , with  $N$  particles of unit mass and periodic boundary conditions. The energy per particle is  $\varepsilon = H_{\text{total}}/N$ . It is not feasible to explore the evolution of finite perturbations of the FPU system at microcanonical equilibrium, due to emerging huge statistical fluctuations. Instead we employed the energy correlation function,  $e(i, t)$  [6,7], which bears the same information as the infinitesimal perturbation in the case of hard-point gas [30]. We performed a massive numerical experiment [25], yet even these efforts were not sufficient to cope with the statistical fluctuations [note the thin green lines in Figs. 4(a) and 4(b)]. Nevertheless, a relatively smooth shape of the ballistic hump allows for a convincing validation of the scaling (5). It is interesting to note that the velocity of the ballistic peaks is determined by the group velocity of effective thermal phonons [7], which therefore can be associated with "walkers" of the CTRW approach.

The perturbation profiles for different values of the energy per particle  $\varepsilon$  reveal another remarkable feature: the central part of profiles is independent of  $\varepsilon$ , while the ballistic humps move as  $\varepsilon$  increases [see Fig. 4(c)]. Such behavior can be derived from the scaling invariance (3). Taking into account that  $K \propto \tau_0^{1-1/\gamma} v_0$ , one can infer that the central part of the LW propagator is invariant under variation of  $v_0$  when  $\tau_0 \propto v_0^{\gamma/(1-\gamma)}$ .

In conclusion, we demonstrated that the collective process of perturbation spreading across two celebrated many-particle systems, a hard-point gas and a Fermi-Pasta-Ulam lattice, is reproduced by a single-particle stochastic process. It is intuitive that an ergodic dynamics is a prerequisite for the diffusionlike perturbation evolution. In integrable




FIG. 4 (color online). (a, b) Energy correlation functions  $e(i, t)$  for the FPU dynamics with  $N = 1.6 \times 10^4$ ,  $\beta = 1$ , and  $\epsilon = 1$  (thin solid lines), compared to the propagators of the generalized LW model with  $\nu_0 = 1.384$  and  $D_\nu = 0.49$  (thick dashed lines). The scaling exponent is  $\gamma = 5/3$ . The FPU profiles have been obtained by averaging over  $5 \times 10^5$  realizations. (c) Energy correlation functions for the FPU system with  $N = 2 \times 10^3$  for the time  $t = 500$  and different values of the energy per particle  $\epsilon$ .

many-body systems, such as a harmonic chain or the Toda lattice [31], the perturbation spreading is a deterministic process, evolving in terms of noninteracting phonons. The task of exploring the sufficient conditions for the CTRW kinetics to occur presents a promising challenge, thereby underpinning the universality of our findings. Thus our results disclose a pathway to explore propagation of information in realistic dissipation-free systems: it allows us to calibrate the transport characteristics of many-body systems (which are beyond the region of validity of the classical Lieb-Robinson theory [11]) by using the parameters of the corresponding random walk process.

Apart from those theoretical challenges, there is room for possible applications. A realization that comes to mind is an array of coupled nanoresonators [32], where a single unit acts as both the receiver and transducer of excitations, which transforms this array into an extended sensor via utilizing the principle of time-of-arrival localization [33].

This work has been supported by the DFG Grants No. HA1517/31-2 (S.D. and P.H.) and No. ZA593/2-1 (V.Z.).

- 
- [1] E. Helfand, *Phys. Rev.* **119**, 1 (1960).  
 [2] A. Torcini, P. Grassberger, and A. Politi, *J. Phys. A* **28**, 4533 (1995); A. Torcini and S. Lepri, *Phys. Rev. E* **55**, R3805 (1997).  
 [3] C. Primo, I.G. Szendro, M.A. Rodriguez, and J.M. Gutierrez, *Phys. Rev. Lett.* **98**, 108501 (2007).  
 [4] G. Giacomelli, R. Hegger, A. Politi, and M. Vassalli, *Phys. Rev. Lett.* **85**, 3616 (2000).  
 [5] P. Cipriani, S. Denisov, and A. Politi, *Phys. Rev. Lett.* **94**, 244301 (2005).

- [6] H. Zhao, *Phys. Rev. Lett.* **96**, 140602 (2006).  
 [7] N. Li, B. Li, and S. Flach, *Phys. Rev. Lett.* **105**, 054102 (2010).  
 [8] J. Dunkel and P. Hänggi, *Phys. Rep.* **471**, 1 (2009); J. Dunkel, P. Talkner, and P. Hänggi, *Phys. Rev. D* **75**, 043001 (2007).  
 [9] C. Marchioro, A. Pellegrinotti, M. Pulvirenti, and L. Triolo, *J. Stat. Phys.* **19**, 499 (1978); P. Butta *et al.*, *J. Stat. Phys.* **127**, 313 (2007).  
 [10] The term “almost” underpins the fact that outside the cone the perturbation is strongly diminished.  
 [11] B. Nachtergaele, H. Raz, B. Schlein, and R. Sims, *Commun. Math. Phys.* **286**, 1073 (2009).  
 [12] E. W. Montroll and G. W. Weiss, *J. Math. Phys. (N.Y.)* **6**, 167 (1965).  
 [13] E. Scalas, in *The Complex Networks of Economic Interactions*, edited by A. Namatame, T. Kaizouji, and Y. Aruka, Lect. Notes Econ. Math. Sys. Vol. 567 (Springer-Verlag, Berlin, 2006), p. 3.  
 [14] E. Barkai, R. Silbey, and G. Zumofen, *Phys. Rev. Lett.* **84**, 5339 (2000).  
 [15] H. Scher and M. Lax, *Phys. Rev. B* **7**, 4491 (1973).  
 [16] A. Blumen, G. Zumofen, and J. Klafter, *Phys. Rev. A* **40**, 3964 (1989).  
 [17] M. F. Shlesinger, G. M. Zaslavsky, and J. Klafter, *Nature (London)* **363**, 31 (1993).  
 [18] J. Klafter and G. Zumofen, *Physica (Amsterdam)* **196A**, 102 (1993).  
 [19] S. Denisov, J. Klafter, and M. Urbakh, *Phys. Rev. Lett.* **91**, 194301 (2003).  
 [20] T. H. Solomon, E. R. Weeks, and H. L. Swinney, *Phys. Rev. Lett.* **71**, 3975 (1993).  
 [21] S. Marksteiner, K. Ellinger, and P. Zoller, *Phys. Rev. A* **53**, 3409 (1996).  
 [22] I. Karatsas and S. Shreve, *Brownian Motion and Stochastic Calculus* (Springer, New York, 1997).  
 [23] This setup amounts to equilibrated initial conditions, when at time  $t = 0$  the particle is in the flight with probability one. It should be distinguished from nonequilibrated conditions when at the time  $t = 0$  the particle is set at rest [15].  
 [24] V. Ziburdaev, M. Schmiedeberg, and H. Stark, *Phys. Rev. E* **78**, 011119 (2008).  
 [25] See supplemental material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.106.180601> for technical details.  
 [26] This quantity can be computed as follows: at time  $t = 0$  the system is locally perturbed,  $\{x_0^p(0) = x_0(0) + \epsilon_x, p_0^p(0) = p_0(0) + \epsilon_p\}$ . Then the two systems, original and perturbed, evolve in parallel. The local excess of energy is given by  $\Delta E(i, t) = H_i^p(t) - H_i(t)$ .  
 [27] Practically, this implies the existence of a time scale,  $t_{LW}$ , beyond which all relevant scattering mechanisms unfold themselves. This time  $t_{LW}$  is system specific.  
 [28] G. Casati and J. Ford, *J. Comput. Phys.* **20**, 97 (1976).  
 [29] J. Ford, *Phys. Rep.* **213**, 271 (1992).  
 [30] L. Delfini *et al.*, *Eur. Phys. J. Special Topics* **146**, 21 (2007).  
 [31] M. Toda, *Nonlinear Waves and Solitons* (Springer, New York, 1989).  
 [32] M. Blencowe, *Phys. Rep.* **395**, 159 (2004).  
 [33] S. Haykin, *Array Signal Processing* (Prentice-Hall, Englewood Cliffs, NJ, 1985).