

```
<?xml version="1.0" encoding="UTF-8"?>
<!DOCTYPE simulation [
<!ENTITY Npts      "64">
<!ENTITY Nsamples  "64">
<!ENTITY L         "3e-5">
]>
<simulation xmds-version="2">
  <name>GPE_1D_spinor_course</name>

  <author> Sebastian Wuester </author>
  <description>
    Gross-Pitaevskii-equation in 1D, SI units
  </description>

  <geometry>
    <propagation_dimension> t </propagation_dimension>
    <transverse_dimensions>
      <dimension name="x" lattice="&Npts;" domain="(-&L;, &L;)" />
      <dimension name="n" type="integer" lattice="3" domain="(0,2)" aliases="k"/>
      <!-- Assignment of components
        0: mF= -1
        1: mF=  0
        2: mF= +1 -->
    </transverse_dimensions>
  </geometry>

  <features>
    <benchmark />
    <auto_vectorise />
    <fftw />
    <globals>
      <![CDATA[
const double hbar = 1.05457266e-34;
const double omega = 10.0*(2.0*M_PI);
const double omega_perp = 200.0*(2.0*M_PI);

// Rb 87
const double mass = 1.4432e-25;
const double a0 = 5.5e-9;
const double a2 = 5.5e-9/101.8*100.4;

//dervied quantities
const double Natoms = 100.0;

const double Us = 4.0*M_PI*hbar*hbar*(a0 + 2.0*a2)/3.0/mass;
const double Ua = 4.0*M_PI*hbar*hbar*(a2 - a0)/3.0/mass;

const double sigmascale = 2.3;
const double sigma = sigmascale*sqrt(hbar/mass/omega);
const double sigma_perp = sqrt(hbar/mass/omega_perp);
const double Ulds = Us/(2.0*M_PI*sigma_perp*sigma_perp);
const double Ulda = Ua/(2.0*M_PI*sigma_perp*sigma_perp);

const double normfact = pow(M_PI*sigma*sigma,-0.25);

]]>
    </globals>
  </features>

  <vector name="wavefunction" initial_space="x n" type="complex">
    <components>psi</components>
```

```
<initialisation>
  <![CDATA[
    double coeff = 0.0;

    if(n == 0)
      coeff = sqrt(0.1);
    if(n == 1)
      coeff = sqrt(0.8);
    if(n == 2)
      coeff = sqrt(0.1);

    psi = normfact*coeff*sqrt(Natoms)*exp(-0.5*x*x/sigma/sigma);
  ]]>
</initialisation>
</vector>

<vector name="potentials" initial_space="x n" type="real">
  <components>trap</components>
  <initialisation>
    <![CDATA[
      trap=0.5*mass*omega*omega*x*x;
    ]]>
  </initialisation>
</vector>

<computed_vector name="auxiliary" dimensions="x" type="real">
  <components> totdens </components>
  <evaluation>
    <dependencies basis="x n"> wavefunction </dependencies>
    <![CDATA[
      totdens = mod2(psi);
    ]]>
  </evaluation>
</computed_vector>

<vector name="coupling" type="real" dimensions="n k">
  <components> F1 </components>
  <initialisation>
    <![CDATA[
      F1 = 0.0;

      // dpsi(-1)/dt
      if(n==0 && (k==0 || k==1))
        F1 = 1.0;
      if(n==0 && k ==2)
        F1 = -1.0;

      // dpsi(0)/dt
      if(n==1 && (k==0 || k==2))
        F1 = 1.0;

      // dpsi(+1)/dt
      if(n==2 && (k==1 || k==2))
        F1 = 1.0;
      if(n==2 && k ==0)
        F1 = -1.0;

    ]]>
  </initialisation>
</vector>
```

```
<computed_vector name="auxiliary2" dimensions="x n" type="real">
  <components> asymmdens </components>
  <evaluation>
    <dependencies basis="x n k"> wavefunction coupling</dependencies>
    <![CDATA[
      asymmdens = F1*mod2(psi( n=>k ));
    ]]>
  </evaluation>
</computed_vector>

<sequence>
  <integrate algorithm="ARK89" interval="4" tolerance="1e-8">
    <samples>200 200 200</samples>
    <operators>
      <operator kind="ip">
        <operator_names>L</operator_names>
        <![CDATA[
          L = -i*0.5*hbar*kx*kx/mass;
        ]]>
      </operator>
      <integration_vectors>wavefunction</integration_vectors>
      <dependencies>potentials auxiliary auxiliary2</dependencies>
      <![CDATA[
        complex conversion = 0.0;

        if(n == 0)
          conversion = conj(psi(n=>2))*psi(n=>1)*psi(n=>1);
        if(n == 1)
          conversion = 2.0*conj(psi)*psi(n=>n+1)*psi(n=>n-1);
        if(n == 2)
          conversion = conj(psi(n=>0))*psi(n=>1)*psi(n=>1);

        dpside_dt = L[psi] - i*( (Ulds*totdens + trap )*psi
          + Ulda*(asymmdens*psi + conversion) )/hbar;

      ]]>
    </operators>
  </integrate>
</sequence>

<output format="hdf5">
  <group>
    <sampling basis="x(&Nsamples;) n" initial_sample="yes">
      <moments>density psire psiim </moments>
      <dependencies>wavefunction </dependencies>
      <![CDATA[
        density = mod2(psi);
        psire = psi.Re();
        psiim = psi.Im();
      ]]>
    </sampling>
  </group>
  <group>
    <sampling basis="kx(&Npts;) n" initial_sample="yes">
      <moments>fspec</moments>
      <dependencies>wavefunction</dependencies>
      <![CDATA[
        fspec = mod2(psi);
      ]]>
    </sampling>
  </group>
</output>
```

```
</sampling>
</group>
<group>
  <sampling basis="kx(0) n" initial_sample="yes">
    <moments>population</moments>
    <dependencies>wavefunction</dependencies>
    <![CDATA[
      population = mod2(psi);
    ]]>
  </sampling>
</group>
</output>
</simulation>
```